



RLP
Approved

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



SOREXA[®] PRO RODENTICIDE GEL

ACTIVE CONSTITUENT: 0.05 g/kg DIFENACOUM

FOR THE CONTROL OF BROWN OR NORWAY RAT (*Rattus norvegicus*) AND MICE, including those resistant to other anticoagulants, in and around industrial, commercial, agricultural, public services and domestic buildings.

A PROFESSIONAL PEST CONTROL PRODUCT

NET CONTENTS: 300 g

BASF Australia Ltd ABN 62 008 437 867.
Level 12, 28 Freshwater Place, Southbank, VIC, 3006, Australia

® Sorexá is a registered trademark of BASF
® Bitrex is a registered trademark of Macfarlan Smith Limited

APVMA Approval No.: 61848/51053

DIRECTIONS FOR USE**RESTRAINTS**

DO NOT place gel in the open.

DO NOT place gel in locations accessible to children and pets.

PEST	SITUATION	RATE	CRITICAL COMMENTS
Mice & Brown or Norway Rats (<i>Rattus norvegicus</i>)	For use in and around (within 2 metres) industrial, commercial, public services, domestic, agricultural buildings (machinery sheds etc).	For mice: use up to 10 g (a bead 10 cm long by 1 cm diameter) per bait point. For rats: use up to 50 g per bait point.	Explore the site and locate runs, nests, burrows, droppings and signs of damage. Eliminate as far as practical, all alternative food sources. Place bait points wherever droppings, damage or other signs of activity are seen. As mice are sporadic feeders use many bait points, not more than 3 metres apart. For rats bait points should be not more than 9 m apart. Use weather resistant bait stations when using bait outdoors.
<p>General information and baiting strategy for rats and mice: Eliminate as far as practical all other food sources. Always place bait where it is inaccessible to children, animals and birds. Protect and cover bait. When using outdoors, use weather resistant bait stations. It is recommended that lockable, securable, weather resistant bait stations such as Roguard be used where possible. Alternatively place bait in suitable covered containers or use weather resistant bait stations constructed from open-ended containers, such as impermeable cardboard boxes or plastic bait stations. Place baits stations under cupboards, behind furniture etc. Try not to touch the bait as this deters mice and rats from eating it. Do not use where food, feed or water could become contaminated. Protect from frost. Place baits only in situations where there is no ready access to occupants of buildings.</p>			<p>INSPECTION: Record the number and locations of bait stations for the purpose of replacement and later removal. Inspect baits frequently, particularly in the first ten days. Replace eaten bait. If the bait is completely consumed, the quantity should be increased. Continue replenishing baits until they are left untouched. Baiting for at least 2 weeks may be necessary to reduce rat/mouse numbers. Most deaths occur 4 to 7 days after consumption of bait, but deaths may occur for up to 2 weeks. Following treatment remove all remains of bait and bait containers and look for and collect dead rats or mice. Burn or bury carcasses. Do not place in refuse bins. DO NOT let cats, dogs or nontarget animals eat poisoned mice and rats. While handling remaining bait material/animal carcasses, wear rubber gloves.</p>

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

GENERAL INSTRUCTIONS

Sorexax® Pro Rodenticide Gel is a very palatable non-spill cereal based all-purpose rodenticidal gel containing the advanced anticoagulant, difenacoum.

The product must only be applied using a caulking gun. Cut off the domed end of the tube taking care not to damage the screw thread. Cut the tapered nozzle to give an opening of approximately 1 cm diameter and screw the nozzle onto the tube. Cover the nozzle when not in use. Sorexax® Pro Rodenticide Gel contains Bitrex® (a human taste deterrent) to help prevent accidental human consumption.

PROTECTION OF LIVESTOCK, WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

The bait is hazardous to pigs, cats, dogs, poultry, birds, and wildlife and care should be exercised in its use.

DO NOT place baits in locations that are accessible to domestic animals, livestock, birds or other animals. DO NOT contaminate dams, waterways and drains with bait or its used containers. DO NOT use this product to control native animals. The product is restricted to use in and around buildings (within 2 metres) or within enclosed spaces. When using outdoors, use weather resistant bait stations. In areas where possums and other nontarget animals frequent, baits should be secured inside proper bait stations or bait stations constructed from drainpipes or boxes.

STORAGE AND DISPOSAL

Store in original container, tightly closed, in a safe place. Triple or preferably pressure rinse containers before disposal. Dispose of rinsings and unused or untaken bait in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit. Empty containers and product should not be burnt.

Search for dead mice and rats – burn or bury them; do not place in refuse bins or rubbish tips. Inspect all bait stations regularly and ensure baits are removed when rodent activity has ceased. Containers that have held bait should not be used for any other purpose.

SAFETY DIRECTIONS

Poisonous if swallowed. May irritate the eyes. Repeated minor exposure may have a cumulative poisoning effect. Avoid contact with eyes. Vitamin K1 (phytomenadione) is antidotal. Wash hands after use.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone 13 11 26. Vitamin K1 (phytomenadione) is antidotal.

MSDS

Additional information is listed in the Material Safety Data Sheet.

NOTE TO PHYSICIANS AND VETERINARIANS

Difenacoum, the active constituent of Sorexax® Pro Rodenticide Gel, is a long lasting anticoagulant chemical, which if ingested by humans, domestic animals or pets, can reduce the clotting power of the blood and haemorrhage may result. Vitamin K1 (phytomenadione) should be administered urgently by medical, appropriate para-medical personnel or veterinarians. If administered intravenously the injection must be given slowly.

MANUFACTURER'S WARRANTY AND EXCLUSION OF LIABILITY

This product as supplied is of a high grade and believed to be suitable for any purpose for which it is expressly recommended and must be used in accordance with the Directions for Use given on the label. No responsibility is accepted in respect of the product save those non-excludable conditions implied by any Federal and State legislation or law of a Territory.

® Sorexax is a registered trademark of BASF

® Bitrex is a registered trademark of Macfarlan Smith Limited

APVMA Approval No.: 61848/51053

Batch No:

Date of Manufacture:



The Chemical Company

BASF Australia Ltd

ABN 62 008 437 867

Level 12, 28 Freshwater Place,

Southbank VIC, 3006, Australia

Website: www.termidor.com.au

Customer Service Hotline: 1800 006 393